

# **APCS Rabbit pAb**

货号: **B17897** 

## 产品信息

反应	Human Maura Dat
区区	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:100
理论分子量	25kDa
实测分子量	25KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,Mouse kidney,Rat liver
细胞定位	Secreted
纯化	Affinity purification


### 抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 100-200 of human APCS (NP_001630 .1).
序列	TSKVIEKFPAPVHICVSWESSSGIAEFWINGTPLVKKGLRQGYFVEAQPKIVLGQEQDSYGGKFDRSQSFVGEIGDLYMW DSVLPPENILSAYQGTPLPAN

靶点信息

研究背景	The protein encoded by this gene is a glycoprotein, belonging to the pentraxin family of proteins, which h as a characteristic pentameric organization. These family members have considerable sequence homolog y which is thought to be the result of gene duplication. The binding of the encoded protein to proteins in t he pathological amyloid cross-beta fold suggests its possible role as a chaperone. This protein is also tho ught to control the degradation of chromatin. It has been demonstrated that this protein binds to apoptoti c cells at an early stage, which raises the possibility that it is involved in dealing with apoptotic cells in viv o.
基因ID	325
基因名	APCS
Swiss	P02743
别名	APCS;HEL-S-92n;PTX2;SAP

### 产品验证



Western blot analysis of APCS expressed in 293T,Mouse kidney,Rat liver using APCS Rabbit pAb at 1:100 0. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Bloc king buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

### 实验步骤

访问官网浏览详情: www.ablybio.cn