

MED1 Rabbit pAb

货号: **B17992**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	61kDa/168kDa
实测分子量	250KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7
细胞定位	Nucleus
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-300 of human MED1 (NP_004765.2).
序列	MKAQGETEESEKLSKMSSLLERLHAKFNQNRPWSETIKLVRQVMEKRVVMSSGGHQHLVSCLETLQKALKVTSLPAMTD RLESIARQNGLGSHLSASGTECYITSDMFYVEVQLDPAGQLCDVKVAHHGENPVSCPELVQQLREKNFDEFKHLKGLVN LYNLPGDNKLKTKMYLALQSLEQDLSKMAIMYWKATNAGPLDKILHGSGVGYLTPRSGGHLMNLYVSPSDLLDDKTASP IILHENNVSRSLGMNASVTIEGTSVYKLPAPLIMGSHPVNDKWTPSFSSITSANSVDLP

靶点信息

研究背景	The activation of gene transcription is a multistep process that is triggered by factors that recognize transcriptional enhancer sites in DNA. These factors work with co-activators to direct transcriptional initiation by the RNA polymerase II apparatus. The protein encoded by this gene is a subunit of the CRSP (cofactor required for SP1 activation) complex, which, along with TFIID, is required for efficient activation by SP1. This protein is also a component of other multisubunit complexes e.g. thyroid hormone receptor-(TR-) associated proteins which interact with TR and facilitate TR function on DNA templates in conjunction with initiation factors and cofactors. It also regulates p53-dependent apoptosis and it is essential for adipogenesis. This protein is known to have the ability to self-oligomerize.
基因ID	5469
基因名	MED1
Swiss	Q15648
别名	MED1;CRSP1;CRSP200;DRIP205;DRIP230;PBP;PPARBP;PPARGBP;RB18A;TRAP220;TRIP2

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn