

Daxx Rabbit pAb

货号: B18011

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	73kDa/75kDa/76kDa/78kDa/81kDa
实测分子量	110KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	K-562,SH-SY5Y
细胞定位	Chromosome,Cytoplasm,Nucleus,PML body,centromere,nucleolus,nucleoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 175-400 of human Daxx (NP_001135441.1).
序列	ASQSPRTRGSRRQIQRLQQLLALYVAEIRRLQEKELDLSELDDPD SAYLQEARLK RKLIRLFGR LCELKDCSSLTGRVIEQRIPYRGTRYPEVNRRIERLINKPGPDTFPDYGDVLR AVEKAAARHSLGLPRQQLQLMAQDAFRDVGIRLQERRHLDLIYNFGCHLTDDYRPGVDPALSDPVLARRLRENRLAMSRLDEVISKYAMLQDKSEEGERKKRRARLQG

靶点信息

研究背景	This gene encodes a multifunctional protein that resides in multiple locations in the nucleus and in the cytoplasm. It interacts with a wide variety of proteins, such as apoptosis antigen Fas, centromere protein C, and transcription factor erythroblastosis virus E26 oncogene homolog 1. In the nucleus, the encoded protein functions as a potent transcription repressor that binds to sumoylated transcription factors. Its repression can be relieved by the sequestration of this protein into promyelocytic leukemia nuclear bodies or nucleoli. This protein also associates with centromeres in G2 phase. In the cytoplasm, the encoded protein may function to regulate apoptosis. The subcellular localization and function of this protein are modulated by post-translational modifications, including sumoylation, phosphorylation and polyubiquitination. Alternative splicing results in multiple transcript variants.
基因ID	1616
基因名	DAXX
Swiss	Q9UER7
别名	DAXX;BING2;DAP6;EAP1

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn