

GET4 Rabbit pAb

货号: **B18029**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	30kDa/36kDa
实测分子量	36kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SKOV3,A-549,HeLa,293T,Mouse brain,Rat lung
细胞定位	Cytoplasm,cytosol
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-327 of human GET4 (NP_057033.2).
序列	MAAAAAMAEQESARNGGRNRGGVQRVEGKLRASVEKGDYYEAHQMYRTLFFRYMSQSKHTEARELMYSGALLFFSHG QQNSAADLSMLVLESLEKAEVEVADELLENLAKVFSLMDPNSPERVTFVSRALKWSSGGSGKLGHPRLHQLLALTLWKEQ NYCESRYHFLHSADGEGCANMLVEYSTSRGFRSEVDMFVAQAVLQFLCLKNKSSASVVFTTYTQKHPSIEDGPPFVEPLL NFIWFLLLAVDGGKLTVFTVLCEQYQPSLRRDPMYNEYLDRIGQLFFGVPPKQTSSYGGLLGNLLTSLMGSSEQEDGEESP SDGSPIELD

靶点信息

研究背景	As part of a cytosolic protein quality control complex, the BAG6/BAT3 complex, maintains misfolded and h ydrophobic patches-containing proteins in a soluble state and participates in their proper delivery to the endoplasmic reticulum or alternatively can promote their sorting to the proteasome where they undergo degradation. The BAG6/BAT3 complex is involved in the post-translational delivery of tail-anchored/type II transmembrane proteins to the endoplasmic reticulum membrane. Recruited to ribosomes, it interacts with the transmembrane region of newly synthesized tail-anchored proteins and together with SGTA and ASNA1 mediates their delivery to the endoplasmic reticulum. Client proteins that cannot be properly delivered to the endoplasmic reticulum are ubiquitinated and sorted to the proteasome. Similarly, the BAG6/BAT3 complex also functions as a sorting platform for proteins of the secretory pathway that are mislocalized to the cytosol either delivering them to the proteasome for degradation or to the endoplasmic reticulum. The BAG6/BAT3 complex also plays a role in the endoplasmic reticulum-associated degradation (ERAD, a quality control mechanism that eliminates unwanted proteins of the endoplasmic reticulum through their retrotranslocation to the cytosol and their targeting to the proteasome. It maintains these retrotranslocated proteins in an unfolded yet soluble state condition in the cytosol to ensure their proper delivery to the proteasome.
基因 ID	51608
基因名	GET4
Swiss	Q7L5D6
别名	GET4;C7orf20;CEE;CGI-20;TRC35

产品验证



Western blot analysis of GET4 expressed in Mouse brain,Rat lung,SKOV3 using GET4 Rabbit pAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Bl ocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn