

EDARADD Rabbit pAb

货号: B18101

产品信息

| | |
|-------|--|
| 反应 | Human |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:2000 |
| 理论分子量 | 23kDa/24kDa |
| 实测分子量 | 25kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | A-431,MCF7 |
| 细胞定位 | Cytoplasm |
| 纯化 | Affinity purification |


抗原信息

| | |
|------|---|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 1-215 of human EDARA DD (NP_665860.2). |
| 序列 | MGLRTTKQMGRGTKAPGHQEDHMKPEVEDTDPSTLSFNMSDKYPIQDTELPKAEECDTITLNCPRNSDMKNQGEENG FPDSTGDPLEISKDNSCKENCTCSSCLLRAPTISDLLNDQDLLDVIRIKLDPCHPTVKNWRNFASKWGMSYDELCFLEQR PQSPTLEFLLRNSQRTVGQLMELCRLYHRADVEKVLRRWVDEEWPKRERGDPSRHF |

靶点信息

| | |
|-------|--|
| 研究背景 | This gene was identified by its association with ectodermal dysplasia, a genetic disorder characterized by defective development of hair, teeth, and eccrine sweat glands. The protein encoded by this gene is a death domain-containing protein, and is found to interact with EDAR, a death domain receptor known to be required for the development of hair, teeth and other ectodermal derivatives. This protein and EDAR are coexpressed in epithelial cells during the formation of hair follicles and teeth. Through its interaction with EDAR, this protein acts as an adaptor, and links the receptor to downstream signaling pathways. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. |
| 基因ID | 128178 |
| 基因名 | EDARADD |
| Swiss | Q8WWZ3 |
| 别名 | EDARADD;ECTD11A;ECTD11B;ED3;EDA3 |

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn