

TXLNG Rabbit pAb

货号: **B18162**

产品信息

| 反应 | Human, Mouse |
|-------|---|
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:2000 |
| 理论分子量 | 46kDa/60kDa |
| 实测分子量 | 75kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | U-87MG,A-549 |
| 细胞定位 | Cytoplasm,Nucleus membrane,cytosol |
| 纯化 | Affinity purification |

抗原信息


| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 1-140 of human TXLNG (NP_060830.2). |
|------|---|
| 序列 | MATRVEEAARGRGGGAEEATEAGRGGRRRSPRQKFEIGTMEEAGICGLGVKADMLCNSQSNDILQHQGSNCGGTSNKH SLEEDEGSDFITENRNLVSPAYCTQESREEIPGGEARTDPPDGQQDSECNRNKEKTLGKEVL |

靶点信息

| 研究背景 | This gene encodes a member of the taxilin family. The encoded protein binds to the C-terminal coiled-coil region of syntaxin family members 1A, 3A and 4A, and may play a role in intracellular vesicle trafficking. This gene is up-regulated by lipopolysaccharide and the gene product may be involved in cell cycle regulation. The related mouse protein was also shown to inhibit activating transcription factor 4-mediated transcription and thus regulate bone mass accrual. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. |
|------|--|
|------|--|

| 基因ID | 55787 |
|-------|-------------------------------------|
| 基因名 | TXLNG |
| Swiss | Q9NUQ3 |
| 别名 | TXLNG;CXorf15;ELRG;FIAT;LSR5;TXLNGX |

产品验证



Western blot analysis of TXLNG expressed in U-87MG,A-549 using TXLNG Rabbit pAb at 1:1000. Secondar y antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5 % non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn