

CAMP Rabbit pAb

货号**: B18312**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	19kDa
实测分子量	18KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	THP-1
细胞定位	Secreted
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 132-170 of human CAM P (NP_004336.3).
序列	NKRFALLGDFFRKSKEKIGKEFKRIVQRIKDFLRNLVPR

靶点信息

rminal signal peptide containing a cathelin domain and a structurally variable cationic antimicrobial de, which is produced by extracellular proteolysis from the C-terminus. In addition to its antibacteri	研究背景	This gene encodes a member of an antimicrobial peptide family, characterized by a highly conserved N-t rminal signal peptide containing a cathelin domain and a structurally variable cationic antimicrobial pept de, which is produced by extracellular proteolysis from the C-terminus. In addition to its antibacterial, and ifungal, and antiviral activities, the encoded protein functions in cell chemotaxis, immune mediator induction, and inflammatory response regulation.
--	------	---

基因ID	820
基因名	САМР
Swiss	P49913
别名	CAMP;CAP-18;CAP18;CRAMP;FALL-39;FALL39;HSD26;LL37

产品验证



Western blot analysis of CAMP expressed in THP-1 using CAMP Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dr y milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn