

ADK Rabbit pAb

货号: B18672

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:100 - 1:500 IF/ICC: 1:50 - 1:200
理论分子量	34kDa/36kDa/38kDa/40kDa
实测分子量	34kDa/36kDa/38kDa/40kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 200-345 of human ADK (NP_001114.2).
序列	PFISQFYKESLMKVMPLYVDILFGNETEAATFAREQGFETDKIKEIAKKTQALPKMNSKRQRIVIFTQGRDDTIMATESEVTAF AVLDQDQKEIIDTNGAGDAFVGGFLSQLVSDKPLTECIRAGHYAASIIIRRTGCTFPEKPDFH

靶点信息

研究背景	This gene an enzyme which catalyzes the transfer of the gamma-phosphate from ATP to adenosine, there by serving as a regulator of concentrations of both extracellular adenosine and intracellular adenine nucl eotides. Adenosine has widespread effects on the cardiovascular, nervous, respiratory, and immune syste ms and inhibitors of the enzyme could play an important pharmacological role in increasing intravascular adenosine concentrations and acting as anti-inflammatory agents. Multiple transcript variants encoding d ifferent isoforms have been found for this gene.
------	--

基因ID	132
基因名	ADK
Swiss	P55263
别名	ADK;AK

产品验证



Western blot analysis of ADK expressed in human liver cancer,mouse brain using ADK Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn