

# MRPS33 Rabbit pAb

货号: **B18738** 

## 产品信息

反应	Human,Mouse
2-1	
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	12kDa
实测分子量	13kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,293T,Mouse brain,Mouse kidney
细胞定位	Mitochondrion
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-106 of human MRPS3 3 (NP_057155.1).	
序列	MSSLSEYAFRMSRLSARLFGEVTRPTNSKSMKVVKLFSELPLAKKKETYDWYPNHHTYAELMQTLRFLGLYRDEHQDFMD EQKRLKKLRGKEKPKKGEGKRAAKRK	

靶点信息

研究背景	Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a l arge 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribo somes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryo tic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the m itoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy rec ognition by sequence homology. The 28S subunit of the mammalian mitoribosome may play a crucial and characteristic role in translation initiation. This gene encodes a 28S subunit protein that is one of the mor e highly conserved mitochondrial ribosomal proteins among mammals, Drosophila and C. elegans. Splice variants that differ in the 5' UTR have been found for this gene; all variants encode the same protein. Pse udogenes corresponding to this gene are found on chromosomes 1q, 4p, 4q, and 20q
基因 <b>ID</b>	51650
基因名	MRPS33
Swiss	Q9Y291
别名	MRPS33;CGI-139;MRP-S33;PTD003;S33mt

#### 产品验证



Western blot analysis of MRPS33 expressed in Mouse brain,A-549,293T using MRPS33 Rabbit pAb at 1: 1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

#### 实验步骤

访问官网浏览详情: www.ablybio.cn