

# **DKK2 Rabbit pAb**

货号: **B18754** 

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:100 - 1:500
理论分子量	28kDa
实测分子量	30kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,NCI-H460
细胞定位	Secreted
纯化	Affinity purification

## 抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 100 to the C-terminus of human DKK 2 (NP_055236.1).
序列	CMVCRRKKKRCHRDGMCCPSTRCNNGICIPVTESILTPHIPALDGTRHRDRNHGHYSNHDLGWQNLGRPHTKMSHIKGH EGDPCLRSSDCIEGFCCARHFWTKICKPVLHQGEVCTKQRKKGSHGLEIFQRCDCAKGLSCKVWKDATYSSKARLHVCQ KI

## 靶点信息

研究背景	This gene encodes a protein that is a member of the dickkopf family. The secreted protein contains two c ysteine rich regions and is involved in embryonic development through its interactions with the Wnt sign aling pathway. It can act as either an agonist or antagonist of Wnt/beta-catenin signaling, depending on t he cellular context and the presence of the co-factor kremen 2. Activity of this protein is also modulated by binding to the Wnt co-receptor LDL-receptor related protein 6 (LRP6).
------	--

基因ID	27123
基因名	DKK2
Swiss	Q9UBU2
别名	DKK2;DKK-2

## 产品验证



Western blot analysis of DKK2 expressed in A-549,NCI-H460 using DKK2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

### 实验步骤

访问官网浏览详情: www.ablybio.cn