

BMP2 Rabbit pAb

货号: **B18841**

产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	44kDa
实测分子量	53kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Rat ovary
细胞定位	Secreted
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human BMP2 (NP_001191.1).
序列	MVAGTRCLLALLLPQVLLGGAAGLVPELGRRKFAAASSGRPSSQPSDEVLSEFELRLLSMFGLKQRPTPSRDAVVPPYMLD LYRRHSGQPGSPAPDHRLE

靶点信息

研究背景	This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of pro teins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMA D family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate each subunit of the disulfide-linked homodimer, which plays a role in bone and car tilage development. Duplication of a regulatory region downstream of this gene causes a form of brachyd actyly characterized by a malformed index finger and second toe in human patients.
------	--

基因 ID	650
基因名	BMP2
Swiss	P12643
别名	BDA2;BMP2A;BMP2

产品验证



Western blot analysis of BMP2 expressed in Rat ovary using BMP2 Rabbit pAb at 1:1000. Secondary antib ody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn