

INMT Rabbit pAb

货号: B18967

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	28kDa
实测分子量	29kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse liver,Mouse lung
细胞定位	Cytoplasm
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-263 of human INMT (NP_006765.4).
序列	MKGGFTGGDEYQKHFLPRDYLATYYSFDSGSPSPEAEMLKFNLECLHKTFGPGLQGDTLIDIGSGPTIYQVLAACDSFQDI TLSDFDRNREELEKWLKEPGAYDWTPAVKFACELEGNSGRWEEKEKLRAAVKRVLKCDVHLGNPLAPAVLPLADCVL TLLAMECACCSDLDAYRAALCNLASLLKPGGHLTTVTLRLPSYMGKREFSCVALEKEEEVEQAVLDAGFDIEQLLHSPQSYS VTNAANNNGVCFIVARKKPGP


靶点信息

研究背景	N-methylation of endogenous and xenobiotic compounds is a major method by which they are degraded. This gene encodes an enzyme that N-methylates indoles such as tryptamine. Alternative splicing results in multiple transcript variants. Read-through transcription also exists between this gene and the downstream FAM188B (family with sequence similarity 188, member B) gene.
------	--

基因ID	11185
基因名	INMT
Swiss	O95050
别名	INMT;TEMT

产品验证

Western blot analysis of INMT expressed in Mouse liver,Mouse lung using INMT Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



实验步骤

访问官网浏览详情: www.ablybio.cn