

# **RPA3 Rabbit pAb**

货号: **B19085** 

# 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	13kDa
实测分子量	19kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse spleen
细胞定位	Nucleus
纯化	Affinity purification


# 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 6-121 of human RPA3 ( NP_002938.1).
序列	DLPRSRINAGMLAQFIDKPVCFVGRLEKIHPTGKMFILSDGEGKNGTIELMEPLDEEISGIVEVVGRVTAKATILCTSYVQFKE DSHPFDLGLYNEAVKIIHDFPQFYPLGIVQHD

靶点信息

研究背景	As part of the heterotrimeric replication protein A complex (RPA/RP-A, binds and stabilizes single-strande d DNA intermediates that form during DNA replication or upon DNA stress. It prevents their reannealing a nd in parallel, recruits and activates different proteins and complexes involved in DNA metabolism. There by, it plays an essential role both in DNA replication and the cellular response to DNA damage. In the cell ular response to DNA damage, the RPA complex controls DNA repair and DNA damage checkpoint activat ion. Through recruitment of ATRIP activates the ATR kinase a master regulator of the DNA damage response.
基因ID	6119
基因名	RPA3
Swiss	P35244
别名	RPA3;REPA3;RP-A p14;RP-Ap14

### 产品验证



Western blot analysis of RPA3 expressed in Mouse spleen using RPA3 Rabbit pAb at 1:1000. Secondary an tibody: HRP Goat Anti-Rabbit lgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% n on-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

### 实验步骤

访问官网浏览详情: www.ablybio.cn