

# ATR Rabbit pAb

货号: B19119

产品信息

|       |  |
|-------|--|
| 反应    | Human,Mouse  |
| 宿主    | Rabbit   |
| 克隆性   | Polyclonal   |
| 预测反应  |  |
| 应用    | WB   |
| 推荐浓度  | <b>WB:</b> 1:100 - 1:500   |
| 理论分子量 | 294kDa/297kDa/301kDa   |
| 实测分子量 | 300KDa   |
| 形式    | Liquid   |
| 保存条件  | Store at -20°C. Avoid freeze / thaw cycles.<br>Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| 偶联物   | Unconjugated   |
| 阳性对照  | HT-29,Mouse testis   |
| 细胞定位  | Chromosome,Nucleus,PML body  |
| 纯化    | Affinity purification  |


抗原信息

|      |  |
|------|--|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 400-650 of human ATR (NP_001175.2).  |
| 序列   | LKMESMEIIIEEIQCQTQQENLSSNSDGISPKRRRLSSSLNPSKRAPKQTEEIKHVDMNQKSILWSALKQKAESLQISLEYSGL<br>KNPVIEMLEGIAVVLQLTALCTVHCShqnmncrtfkdcqhkskkkpsvvitwmsldfytkvlkscrsllsvqkldleati<br>dkvvkiydaliymqvnssfedhiledlcmllslpwiyshsddgclklttfaanlltlscrisdssyspqaqsrvflltlfpr<br>iflE |

靶点信息

|       |   |
|-------|---|
| 研究背景  | The protein encoded by this gene belongs the PI3/PI4-kinase family, and is most closely related to ATM, a protein kinase encoded by the gene mutated in ataxia telangiectasia. This protein and ATM share similarity with Schizosaccharomyces pombe rad3, a cell cycle checkpoint gene required for cell cycle arrest and DNA damage repair in response to DNA damage. This kinase has been shown to phosphorylate checkpoint kinase CHK1, checkpoint proteins RAD17, and RAD9, as well as tumor suppressor protein BRCA1. Mutations of this gene are associated with Seckel syndrome. An alternatively spliced transcript variant of this gene has been reported, however, its full length nature is not known. Transcript variants utilizing alternative polyA sites exist. |
| 基因ID  | 545   |
| 基因名   | ATR   |
| Swiss | Q13535  |
| 别名    | ATR;FCTCS;FRP1;MEC1;SCKL;SCKL1  |

产品验证



Western blot analysis of ATR expressed in HT-29, Mouse testis using ATR Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)