

HIC1 Rabbit pAb

货号: B19136

产品信息

反应	Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	74kDa/76kDa
实测分子量	76kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse liver
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 180-430 of human HIC1 (NP_001091672.1).
序列	AATPVIQACYPSPVGPPPPAAEPPSGPEAAVNTHCAELYASGPGPAAALCASERRCSPLCGLDLSKKSPPGSAAPERPLAE RELPPRPDSPPSAGPAAYKEPPLALPSLPPLPFQKLEEAAPSDPFRGGSGSPGPEPPGRPDGPSLLYRWMKHEPGLGSYG DELGRERGSPPERCEERGGDAAVSPGGPPLGLAPPPRYPGSLDGPAGGDGDDYKSSEETGSSEDPSPPGGHLEGYP CPHLAYGEPE

靶点信息

研究背景	This gene functions as a growth regulatory and tumor repressor gene. Hypermethylation or deletion of the region of this gene have been associated with tumors and the contiguous-gene syndrome, Miller-Dieker syndrome. Alternative splicing of this gene results in multiple transcript variants.
基因ID	3090

基因名	HIC1
Swiss	Q14526
别名	HIC1;ZBTB29;ZNF901;hic-1

产品验证



Western blot analysis of HIC1 expressed in Mouse liver using HIC1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn