

# FA2H Rabbit pAb

货号: B19159

## 产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	18kDa/42kDa
实测分子量	40kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse brain,Mouse stomach,Mouse kidney,Rat brain
细胞定位	Endoplasmic reticulum membrane,Microsome membrane,Multi-pass membrane protein
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 95-170 of human FA2H ( NP_077282.3).
序列	NEPVALEETQKTDPA MEPRFKVVDWDKDLVDWRKPLLWQVGH LG EKYDEWVHQPVTRPIRLFHSDLIEGLSKTVWY

## 靶点信息

研究背景	This gene encodes a protein that catalyzes the synthesis of 2-hydroxysphingolipids, a subset of sphingolipids that contain 2-hydroxy fatty acids. Sphingolipids play roles in many cellular processes and their structural diversity arises from modification of the hydrophobic ceramide moiety, such as by 2-hydroxylation of the N-acyl chain, and the existence of many different head groups. Mutations in this gene have been associated with leukodystrophy dysmyelinating with spastic paraparesis with or without dystonia.
基因ID	79152

基因名	FA2H
Swiss	Q7L5A8
别名	FA2H;FAAH;FAH1;FAXDC1;SCS7;SPG35

产品验证



Western blot analysis of FA2H expressed in Mouse brain,Rat brain,Mouse stomach using FA2H Rabbit pA b at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)