

URM1 Rabbit pAb

货号: **B19180**

产品信息

| 反应 | Mouse,Rat |
|-------|---|
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB IF/ICC |
| 推荐浓度 | WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:100 |
| 理论分子量 | 7kDa/11kDa/15kDa |
| 实测分子量 | 11kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | Rat eye |
| 细胞定位 | Cytoplasm |
| 纯化 | Affinity purification |

抗原信息


| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 1-63 of human URM1 (N P_001252511.1). |
|------|---|
| 序列 | MAAPLSVEVEFGGGAELLFDGIKKHRVTLPGQEEPWDIRNLLIWIKKNLLKERPELFIQGDSV |

靶点信息

| 研究背景 | Acts as a sulfur carrier required for 2-thiolation of mcm(5S(2U at tRNA wobble positions of cytosolic tRNA(Lys, tRNA(Glu and tRNA(Gln. Serves as sulfur donor in tRNA 2-thiolation reaction by being thiocarboxylate d (-COSH at its C-terminus by MOCS3. The sulfur is then transferred to tRNA to form 2-thiolation of mcm(5S(2U. Also acts as a ubiquitin-like protein (UBL that is covalently conjugated via an isopeptide bond to ly sine residues of target proteins such as MOCS3, ATPBD3, CTU2, USP15 and CAS. The thiocarboxylated for m serves as substrate for conjugation and oxidative stress specifically induces the formation of UBL-prote in conjugates. |
|------|---|
|------|---|

| 基因 ID | 81605 |
|--------------|--------------|
| 基因名 | URM1 |
| Swiss | Q9BTM9 |
| 别名 | URM1;C9orf74 |

产品验证



Western blot analysis of URM1 expressed in Rat eye using URM1 Rabbit pAb at 1:1000. Secondary antibod y: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn