

# FGL2 Rabbit pAb

货号: **B19200** 

### 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	50kDa
实测分子量	70kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2
细胞定位	Secreted
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 24-140 of human FGL2 ( NP_006673.1).	
序列	NNETEEIKDERAKDVCPVRLESRGKCEEAGECPYQVSLPPLTIQLPKQFSRIEEVFKEVQNLKEIVNSLKKSCQDCKLQADD NGDPGRNGLLLPSTGAPGEVGDNRVRELESEVNKL	

### 靶点信息

研究背景	The protein encoded by this gene is a secreted protein that is similar to the beta- and gamma-chains of fi brinogen. The carboxyl-terminus of the encoded protein consists of the fibrinogen-related domains (FRED ). The encoded protein forms a tetrameric complex which is stabilized by interchain disulfide bonds. This protein may play a role in physiologic functions at mucosal sites.
基因 <b>ID</b>	10875

基因名	FGL2
Swiss	Q14314
别名	FGL2;T49;pT49

#### 产品验证



Western blot analysis of FGL2 expressed in HepG2 using FGL2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dr y milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

### 实验步骤

访问官网浏览详情: www.ablybio.cn