

# PP2A-B55 $\alpha$ /PR55 $\alpha$ /PPP2R2A Rabbit pAb

货号: B19295

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	<a href="#">WB</a>
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	51kDa/53kDa
实测分子量	52kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Jurkat,Mouse brain
细胞定位	cytosol,glutamatergic synapse,nucleoplasm
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-447 of human PP2A-B55 $\alpha$ /PR55 $\alpha$ /PPP2R2A (NP_002708.1).
序列	MAGAGGGNDIQWCFSQVKGAVIDDVAEADIISTVEFNHSPELLATGDKGGRVVIFQQEQENKIQSHSRGEYNVYSTFQSHEPEFDYLKSLEIEEKINKIRWLPQKNAAQFLLSTNDKTIKLWKISERDKRPEGYNLKEEDGRYRDPTTVTTLRVPVFRPMDSLVEASPRIFANAHYHINSISINSDYETYLSADDLRINLWHEITDRSFNIVDIK PANMEELTEVITAAEFHPNSCNTFVYSSSKGTIRLCDMRASALCDRHSKLFEEDPSNRSSFEIISISDVKFHSGRYMMTRDYLSQLKIWL DNMENRPVETYQVHEYLRSKLCQLYENDCIFDKFECCWNGSDSVMTGSYNNFFRMFDRNTKRDITLEASRENNKPRTVLKPRKVCASGKRKKDEISVDSLDFNKKILHTAWHPKENIIAVATTNNLYIFQDKVN

## 靶点信息

研究背景	The product of this gene belongs to the phosphatase 2 regulatory subunit B family. Protein phosphatase 2 is one of the four major Ser/Thr phosphatases, and it is implicated in the negative control of cell growth and division. It consists of a common heteromeric core enzyme, which is composed of a catalytic subunit and a constant regulatory subunit, that associates with a variety of regulatory subunits. The B regulatory subunit might modulate substrate selectivity and catalytic activity. This gene encodes an alpha isoform of the regulatory subunit B55 subfamily. Alternatively spliced transcript variants have been described.
基因ID	5520
基因名	PPP2R2A
Swiss	P63151
别名	PPP2R2A;B55A;B55ALPHA;PR52A;PR55A;PR55alpha

## 产品验证



Western blot analysis of PP2A-B55α/PR55α/PPP2R2A expressed in HeLa,Jurkat,Mouse brain using PP2A-B55α/PR55α/PPP2R2A Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit Ig G (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST . Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)