

# MCM4 Rabbit pAb

# 货号**: B19301**

# 产品信息

| 反应    | Human,Mouse,Rat  |
|-------|--|
| 宿主    | Rabbit   |
| 克隆性   | Polyclonal   |
| 预测反应  |  |
| 应用    | WB IHC   |
| 推荐浓度  | WB: 1:500 - 1:2000<br>IHC: 1:100 - 1:200   |
| 理论分子量 | 96kDa  |
| 实测分子量 | 110kDa   |
| 形式    | Liquid   |
| 保存条件  | Store at -20°C. Avoid freeze / thaw cycles.<br>Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物   | Unconjugated   |
| 阳性对照  | HeLa,Jurkat,Raji   |
| 细胞定位  | Nucleus  |
| 纯化    | Affinity purification  |

# 抗原信息

| 抗原信息 | A synthetic peptide corresponding to a sequence within amino acids 350-450 of human MCM4 (NP_87742 3.1).  |  |
|------|---|--|
| 序列   | QMIKLQESPEDMPAGQTPHTVILFAHNDLVDKVQPGDRVNVTGIYRAVPIRVNPRVSNVKSVYKTHIDVIHYRKTDAKRLH<br>GLDEEAEQKLFSEKRVELLK |  |

### 靶点信息

| 研究背景  | The protein encoded by this gene is one of the highly conserved mini-chromosome maintenance proteins (MCM) that are essential for the initiation of eukaryotic genome replication. The hexameric protein compl ex formed by MCM proteins is a key component of the pre-replication complex (pre_RC) and may be invol ved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 6 and 7 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. The phosphorylation of this protein by CDC2 kinase reduces th e DNA helicase activity and chromatin binding of the MCM complex. This gene is mapped to a region on t he chromosome 8 head-to-head next to the PRKDC/DNA-PK, a DNA-activated protein kinase involved in th e repair of DNA double-strand breaks. Alternatively spliced transcript variants encoding the same protein have been reported. |
|-------|--|
| 基因ID  | 4173   |
| 基因名   | MCM4   |
| Swiss | P33991   |
| 别名    | MCM4;CDC21;CDC54;NKCD;NKGCD;P1-CDC21;hCdc21  |

#### 产品验证



Western blot analysis of MCM4 expressed in HeLa,Jurkat,Raji using MCM4 Rabbit pAb at 1:1000. Secon dary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking b uffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

#### 实验步骤

访问官网浏览详情: www.ablybio.cn