

CSNK2A2 Rabbit pAb

货号: B19313

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:100 IF/ICC: 1:50 - 1:100
理论分子量	41kDa
实测分子量	38kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	K-562,Mouse testis
细胞定位	acrosomal vesicle,cytosol,nucleoplasm,nucleus,plasma membrane
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-350 of human CSNK2 A2 (NP_001887.1).
序列	MPGPAAGSRARVYAEVNSLRsREYWDYEAHVPSWGNQDDYQLVRKLGRGKYSEVF E AINITNN E RVVVKILKPVK K KKIK REVKILENLRGGTNI K LIDTVKDPVSKTPALVFEYINNTDFKQLYQILTD F DIRFYMYELLKALDYCHSKGIMHRDVKPHN V MI DHQQKKLRLIDWGLAEFYHPAQEYNVRVASRYFKGPELLVDYQMYDYS L DMWSLGCMLASMIFRREPFFHGQDN Y DQL VRIAKVLGTEELYGYLK K YHIDLDPHFNDILGQHSRKRWENFIHSEN R HLVSPEALDLLDKLLRYDHQ Q RLTAKEAMEHPYF YPVVKEQSQPCADNAVLSSGLTAAR

靶点信息

研究背景	Catalytic subunit of a constitutively active serine/threonine-protein kinase complex that phosphorylates a large number of substrates containing acidic residues C-terminal to the phosphorylated serine or threonine. Regulates numerous cellular processes, such as cell cycle progression, apoptosis and transcription, as well as viral infection. May act as a regulatory node which integrates and coordinates numerous signals leading to an appropriate cellular response. During mitosis, functions as a component of the p53/TP53-dependent spindle assembly checkpoint (SAC) that maintains cyclin-B-CDK1 activity and G2 arrest in response to spindle damage. Also required for p53/TP53-mediated apoptosis, phosphorylating 'Ser-392' of p53/TP53 following UV irradiation.
基因ID	1459
基因名	CSNK2A2
Swiss	P19784
别名	CSNK2A2;CK2A2;CK2alpha;CSNK2A1

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn