

ARMET/ARP/MANF Rabbit pAb

货号: B19353

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	20kDa
实测分子量	16KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HT-29,HepG2,BxPC-3,Mouse testis,Mouse pancreas,Rat testis
细胞定位	Secreted
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 25-182 of human ARME T/ARP/MANF (NP_006001.4).
序列	SRALRPGDCEVCISYLGRFYQDLKDRDVTFSPATIENELIKFCREARGKENRLCYYIGATDDAATKIINEVSKPLAHHIPVEKIC EKLKKKDSQICELKYDKQIDLSTVDLKKLRVKELKKILDDWGETCKGCAEKSDYIRKINELMPKYAPKAASAR

靶点信息

研究背景	The protein encoded by this gene is localized in the endoplasmic reticulum (ER) and golgi, and is also secr eted. Reducing expression of this gene increases susceptibility to ER stress-induced death and results in cell proliferation. Activity of this protein is important in promoting the survival of dopaminergic neurons. The presence of polymorphisms in the N-terminal arginine-rich region, including a specific mutation that changes an ATG start codon to AGG, have been reported in a variety of solid tumors; however, these polymorphisms were later shown to exist in normal tissues and are thus no longer thought to be tumor-related.
基因 ID	7873
基因名	MANF
Swiss	P55145
别名	MANF;ARMET;ARP

产品验证



Western blot analysis of ARMET/ARP/MANF expressed in Mouse testis,Rat testis,HT-29 using ARMET/ARP/MANF Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit lgG (H+L) at 1:5000. Ly sates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhance d Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn