

USP36 Rabbit pAb

货号: **B19473**

产品信息

反应	Human,Mouse
12,127.	numan, Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:100
理论分子量	122kDa
实测分子量	123kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,A-431,HeLa
细胞定位	Nucleus, nucleolus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-120 of human USP36 (NP_079366.3).
序列	MPIVDKLKEALKPGRKDSADDGELGKLLASSAKKVLLQKIEFEPASKSFSYQLEALKSKYVLLNPKTEGASRHKSGDDPPA RRQGSEHTYESCGDGVPAPQKVLFPTERLSLRWERVFRV

靶点信息

研究背景	This gene encodes a member of the peptidase C19 or ubiquitin-specific protease family of cysteine prote ases. Members of this family remove ubiquitin molecules from polyubiquitinated proteins. The encoded protein may deubiquitinate and stabilize the transcription factor c-Myc, also known as MYC, an important on coprotein known to be upregulated in most human cancers. The encoded protease may also regulate the activation of autophagy. This gene exhibits elevated expression in some breast and lung cancers.
	activation of autophagy. This gene exhibits elevated expression in some breast and lung cancers.

基因ID	57602
基因名	USP36
Swiss	Q9P275
别名	USP36;DUB1

产品验证



Western blot analysis of USP36 expressed in U-87MG,A-431,HeLa using USP36 Rabbit pAb at 1:1000. S econdary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocki ng buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn