

MAFF Rabbit pAb

货号: **B19541**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	14kDa/17kDa
实测分子量	18kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HT-29
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 75-164 of human MAFF (NP_036455.1).	
序列	QKEELQKQKSELEREVDKLARENAAMRLELDALRGKCEALQGFARSVAAARGPATLVAPASVITIVKSTPGSGSGPAHGPD PAHGPASCS	

靶点信息

研究背景	The protein encoded by this gene is a basic leucine zipper (bZIP) transcription factor that lacks a transact ivation domain. It is known to bind the US-2 DNA element in the promoter of the oxytocin receptor (OTR) gene and most likely heterodimerizes with other leucine zipper-containing proteins to enhance expression of the OTR gene during term pregnancy. The encoded protein can also form homodimers, and since it I acks a transactivation domain, the homodimer may act as a repressor of transcription. This gene may als o be involved in the cellular stress response. Multiple transcript variants encoding two different isoforms have been found for this gene.
------	--

基因 ID	23764
基因名	MAFF
Swiss	Q9ULX9
别名	MAFF;U-MAF;hMafF

产品验证



Western blot analysis of MAFF expressed in HT-29 using MAFF Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dr y milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn