

ITIH4 Rabbit pAb

货号: **B19593**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	93kDa/99kDa/101kDa/103kDa
实测分子量	103kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	LO2,A-549
细胞定位	Secreted
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 689-930 of human ITIH4 (NP_002209.2).
序列	RLAILPASAPPATSNPDPAVSRVMNMKIEETTMTTQTPAPIQAPSAILPLPGQSVERLCVDPRHRQGPVNLLSDPEQGVEVT GQYEREKAGFSWIEVTFKNPLVWVHASPEHVVVTRNRRSSAYKWKETLFSVMPGLKMTMDKTGLLLLSDPDKVTIGLLF WDGRGEGLRLLLRDTDRFSSHVGGTLGQFYQEVLWGSPAASDDGRRTLRVQGNDHSATRERRLDYQEGPPGVEISCW SVEL

靶点信息

wo smaller forms. Expression of this gene has been detected only in liver, and it seems to be upregulated during surgical trauma. This gene is part of a cluster of similar genes on chromosome 3. Two transcript variants encoding different isoforms have been found for this gene.

基因ID	3700
基因名	ITIH4
Swiss	Q14624
别名	ITIH4;GP120;H4P;IHRP;ITI-HC4;ITIHL1;PK-120;PK120

产品验证



Western blot analysis of ITIH4 expressed in LO2,A-549 using ITIH4 Rabbit pAb at 1:1000. Secondary antib ody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn