

VASH1 Rabbit pAb

货号: B19665

产品信息

| | |
|-------|--|
| 反应 | Human,Mouse |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:2000 |
| 理论分子量 | 21kDa/40kDa |
| 实测分子量 | 41kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | Mouse brain,Mouse stomach |
| 细胞定位 | Secreted |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|--|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 1-365 of human VASH1 (NP_055724.1). |
| 序列 | MPGGKKVAGGGSSGATPTSAAATAPSGVRRLETSEGTSAQRDEEPEEEGEEDLRDGGVPFFNRGGLPVDEATWERM WKHVAKIHPDGEKVAQRIRGATDLPKIPIPSVPTFQPSTPVPERLEAVQRYIRELQYNHTGTQFFEIKKSRLTGLMDLAKE TKEALPIKCLEAVILGIYLTNSMPTLERFPISFKTYFSGNYFRHIVLGVNFAGRYGALGMSRREDLMLYKPPAFRTLSELVLD FE AAYGRCWHVLLKKVKLGQSVSHDPHSVEQIEWKHSVLDVERLGRDDFRKELERHARDMRLKIGKGTGPPSPTKDRKKDV SSPQRQAQSSPHRRNSRSERRPSGDKKTSEPKAMPDLNGYQIRV |

靶点信息

| | |
|-------|---|
| 研究背景 | Tyrosine carboxypeptidase that removes the C-terminal tyrosine residue of alpha-tubulin, thereby regulating microtubule dynamics and function. Critical for spindle function and accurate chromosome segregation during mitosis since microtubule detyronisation regulates mitotic spindle length and positioning. Acts as an angiogenesis inhibitor: inhibits migration, proliferation and network formation by endothelial cells as well as angiogenesis. This inhibitory effect is selective to endothelial cells as it does not affect the migration of smooth muscle cells or fibroblasts. |
| 基因ID | 22846 |
| 基因名 | VASH1 |
| Swiss | Q7L8A9 |
| 别名 | VASH1;KIAA1036 |

产品验证

Western blot analysis of VASH1 expressed in Mouse brain,Mouse stomach using VASH1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



实验步骤

访问官网浏览详情: www.ablybio.cn