

FCGR2B Rabbit pAb

货号: B19671

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	31kDa/33kDa/34kDa
实测分子量	40kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse spleen,Rat liver
细胞定位	Cell membrane,Single-pass type I membrane protein
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 43-217 of human FCGR2B (P31994).
序列	TPAAPPKAVLKEPQWINVLQEDSVTLCRGTHSPESDSIQWFHNGNLIPTHTQPSYRFKANNNDSGEYTCQTGQTSLSDPVHLTVLSEWLVLQTPHLEFQEGETIVLRCHSWDKPLVKVTFFQNGKSKFSRSDPNFSIPQANHSHGSDYHCTGNIGYTLYSSKPVTITVQAP


靶点信息

研究背景	The protein encoded by this gene is a low affinity receptor for the Fc region of immunoglobulin gamma complexes. The encoded protein is involved in the phagocytosis of immune complexes and in the regulation of antibody production by B-cells. Variations in this gene may increase susceptibility to systemic lupus erythematosus (SLE). Several transcript variants encoding different isoforms have been found for this gene.
------	---

基因ID	2213
基因名	FCGR2B
Swiss	P31994
别名	FCGR2B;CD32;CD32B;FCG2;FCGR2;IGFR2

产品验证

Western blot analysis of FCGR2B expressed in Mouse spleen,Rat liver using FCGR2B Rabbit pAb at 1:1000 . Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



实验步骤

访问官网浏览详情: www.ablybio.cn