

# COX6B1 Rabbit pAb

货号: **B19716** 

# 产品信息

Mouse,Rat
Rabbit
Polyclonal
WB IHC
WB: 1:500 - 1:2000 IHC: 1:100 - 1:200
10kDa
12kDa
Liquid
Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Unconjugated
Mouse brain, Mouse liver, Rat liver
Mitochondrion intermembrane space
Affinity purification


# 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-86 of human COX6B1 (NP_001854.1).	
序列	MAEDMETKIKNYKTAPFDSRFPNQNQTRNCWQNYLDFHRCQKAMTAKGGDISVCEWYQRVYQSLCPTSWVTDWDEQ RAEGTFPGKI	

靶点信息

研究背景	Cytochrome c oxidase (COX), the terminal enzyme of the mitochondrial respiratory chain, catalyzes the el ectron transfer from reduced cytochrome c to oxygen. It is a heteromeric complex consisting of 3 catalyti c subunits encoded by mitochondrial genes and multiple structural subunits encoded by nuclear genes. T he mitochondrially-encoded subunits function in electron transfer, and the nuclear-encoded subunits may be involved in the regulation and assembly of the complex. This nuclear gene encodes subunit VIb. Mutati ons in this gene are associated with severe infantile encephalomyopathy. Three pseudogenes COX6BP-1, COX6BP-2 and COX6BP-3 have been found on chromosomes 7, 17 and 22q13.1-13.2, respectively.
基因ID	1340
基因名	COX6B1
Swiss	P14854
别名	COX6B1;COX6B;COXG;COXVIb1

### 产品验证



Western blot analysis of COX6B1 expressed in Mouse brain, Mouse liver, Rat liver using COX6B1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 12 0s.

### 实验步骤

访问官网浏览详情: www.ablybio.cn