

IKZF2 Rabbit pAb

货号: **B19767**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	5kDa/16kDa/25kDa/26kDa/51kDa/56kDa/59kDa
实测分子量	60kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	K-562,HeLa,Mouse thymus,Rat thymus
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信	恴	Recombinant fusion protein containing a sequence corresponding to amino acids 300-440 of human IKZF 2 (NP_001072994.1).
序列	IJ	TYLGAEALHPLMQHPPSTIAEVAPVISSAYSQVYHPNRIERPISRETADSHENNMDGPISLIRPKSRPQEREASPSNSCLDST DSESSHDDHQSYQGHPALNPKRKQSPAYMKEDVKALDTTKAPKGSLKDIYKVFNGEGE

靶点信息

研究背景	This gene encodes a member of the Ikaros family of zinc-finger proteins. Three members of this protein fa mily (Ikaros, Aiolos and Helios) are hematopoietic-specific transcription factors involved in the regulation of lymphocyte development. This protein forms homo- or hetero-dimers with other Ikaros family members , and is thought to function predominantly in early hematopoietic development. Multiple transcript varian ts encoding different isoforms have been found for this gene, but the biological validity of some variants has not been determined.
------	--

基因 ID	22807
基因名	IKZF2
Swiss	Q9UKS7
别名	IKZF2;ANF1A2;HELIOS;ZNF1A2;ZNFN1A2

产品验证



Western blot analysis of IKZF2 expressed in Mouse thymus,Rat thymus,K-562 using IKZF2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane . Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn