

# CHD9 Rabbit pAb

货号: B19815

产品信息

|       |  |
|-------|--|
| 反应    | Human,Mouse  |
| 宿主    | Rabbit   |
| 克隆性   | Polyclonal   |
| 预测反应  |  |
| 应用    | WB   |
| 推荐浓度  | <b>WB:</b> 1:500 - 1:2000  |
| 理论分子量 | 324kDa/326kDa  |
| 实测分子量 | 300kDa   |
| 形式    | Liquid   |
| 保存条件  | Store at -20°C. Avoid freeze / thaw cycles.<br>Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物   | Unconjugated   |
| 阳性对照  | U-87MG,HeLa,Jurkat,A-549,293T,K-562,Mouse kidney   |
| 细胞定位  | Cytoplasm,Nucleus  |
| 纯化    | Affinity purification  |

抗原信息


|      |   |
|------|---|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 85-190 of human CHD9 (NP_079410.4). |
| 序列   | SFGSPAEHVLSPHSQFNCSPHPQNQPNGLFPDVSDGSPMWDGHTATTISNQNGSPFHQQGHSHSMHQNKSFVAHHD<br>FALFQANEQQTQCTSLRSQQNRNNLNPGQ     |

靶点信息

|      |  |
|------|--|
| 研究背景 | Acts as a transcriptional coactivator for PPARA and possibly other nuclear receptors. Proposed to be a AT P-dependent chromatin remodeling protein. Has DNA-dependent ATPase activity and binds to A/T-rich DN A. Associates with A/T-rich regulatory regions in promoters of genes that participate in the differentiation of progenitors during osteogenesis (By similarity. |
| 基因ID | 80205  |

|       |                                      |
|-------|--------------------------------------|
| 基因名   | CHD9                                 |
| Swiss | Q3L8U1                               |
| 别名    | CHD9;AD013;CHD-9;CReMM;KISH2;PRIC320 |

产品验证



Western blot analysis of CHD9 expressed in Mouse kidney,U-87MG,HeLa using CHD9 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)