

NACC1 Rabbit pAb

货号: B19912

产品信息

| | |
|-------|--|
| 反应 | Human,Mouse |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB IHC |
| 推荐浓度 | WB: 1:500 - 1:2000 IHC: 1:100 - 1:200 |
| 理论分子量 | 57kDa |
| 实测分子量 | 57kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | SKOV3,Hela,Jurkat,Mouse brain,Mouse kidney |
| 细胞定位 | Cytoplasm,Nucleus |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|--|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 170-340 of human NAC C1 (NP_443108.1). |
| 序列 | QQESDSVQCMPVAKRLWDSGQKEAGGGGNGSRKMAKFSTPDLAANRPHQPPPPQAPVVAAAQPAVAAGAGQPAGG VAAAGGVVSGPSTERTSPGTSSAYTSDSPGSYHNEEDEEDGGEEGMDEQYRQICNMYTMYSMMNVGQTAEKVEALP EQVAPESRNRIRVRQDL |

靶点信息

| | |
|------|--|
| 研究背景 | This gene encodes a member of the BTB/POZ protein family. BTB/POZ proteins are involved in several cell ular processes including proliferation, apoptosis and transcription regulation. The encoded protein is a tra nscriptional repressor that plays a role in stem cell self-renewal and pluripotency maintenance. The enco ded protein also suppresses transcription of the candidate tumor suppressor Gadd45GIP1, and expression of this gene may play a role in the progression of multiple types of cancer. A pseudogene of this gene is l ocated on the short arm of chromosome 9. |
|------|--|

| | |
|-------|---|
| 基因ID | 112939 |
| 基因名 | NACC1 |
| Swiss | Q96RE7 |
| 别名 | NACC1;BEND8;BTBD14B;BTBD30;NAC-1;NAC1;NECFM |

产品验证



Western blot analysis of NACC1 expressed in Mouse brain,SKOV3,Hela using NACC1 Rabbit pAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Bl ocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn