

OCIAD1 Rabbit pAb

货号: B19938

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	20kDa/21kDa/27kDa
实测分子量	28kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse liver
细胞定位	Endosome
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-245 of human OCIAD1 (NP_060300.1).
序列	MNGRADFREPNAEVPRPIPHIGPDYIPTEEERRVFAECNDESFWFRSVPLAATSMELITQGLISKGILSSHPKYGSIPKLILACIM GYFAGKLSYVKTCQEKFKKLENSPLGEALRSGQARRSSPPGHYYQKSKYDSSVSGQSSFVTSPAADNIEMLPHYEPIPFSS SMNESAPTGITDHIVQGPDPNLEESPKRKNITYEELRNKNRESYEVSLTQKTDPSVRPMHERVPKKEVKVNKYGDTWDE

靶点信息

研究背景	Maintains stem cell potency (By similarity. Increases STAT3 phosphorylation and controls ERK phosphorylation (By similarity. May act as a scaffold, increasing STAT3 recruitment onto endosomes (By similarity. Involved in integrin-mediated cancer cell adhesion and colony formation in ovarian cancer.
基因ID	54940

基因名	OCIAD1
Swiss	Q9NX40
别名	OCIAD1;ASRIJ;OCIA;TPA018

产品验证



Western blot analysis of OCIAD1 expressed in Mouse liver using OCIAD1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn