

BCL3 Rabbit pAb

货号: **B19953**

产品信息

| 反应 | Human,Mouse,Rat |
|-------|---|
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB IF/ICC |
| 推荐浓度 | WB: 1:500 - 1:1000 IF/ICC: 1:50 - 1:200 |
| 理论分子量 | 47kDa |
| 实测分子量 | 60kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | Raji,293T |
| 细胞定位 | Cytoplasm,Nucleus,perinuclear region |
| 纯化 | Affinity purification |

抗原信息


| 抗原信息 | A synthetic peptide corresponding to a sequence within amino acids 300-400 of human BCL3 (NP_005169 .2). |
|------|---|
| 序列 | ANVNAQMYSGSSALHSASGRGLLPLVRTLVRSGADSSLKNCHNDTPLMVARSRRVIDILRGKATRPASTSQPDPSPDRSA NTSPESSSRLSSNGLLSASPS |

靶点信息

| 研究背景 | This gene is a proto-oncogene candidate. It is identified by its translocation into the immunoglobulin alph a-locus in some cases of B-cell leukemia. The protein encoded by this gene contains seven ankyrin repeat s, which are most closely related to those found in I kappa B proteins. This protein functions as a transcrip tional co-activator that activates through its association with NF-kappa B homodimers. The expression of this gene can be induced by NF-kappa B, which forms a part of the autoregulatory loop that controls the n uclear residence of p50 NF-kappa B. |
|------|---|
|------|---|

| 基因ID | 602 |
|-------|------------------|
| 基因名 | BCL3 |
| Swiss | P20749 |
| 别名 | BCL3;BCL4;D19S37 |

产品验证



Western blot analysis of BCL3 expressed in Raji,293T using BCL3 Rabbit pAb at 1:1000. Secondary antibo dy: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-f at dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn