

TNFRSF10C Rabbit pAb

货号**: B19973**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	27kDa
实测分子量	27KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse liver,Rat liver
细胞定位	Cell membrane,GPI-anchor,Lipid-anchor
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 150-238 of human TNFR SF10C (NP_003832.2).	
序列	VEEFGANATVETPAAEETMNTSPGTPAPAAEETMNTSPGTPAPAAEETMTTSPGTPAPAAEETMTTSPGTPAPAAEETMTT SPGTPASS	

靶点信息

研究背景	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor contains an extracellular TRAIL-binding domain and a transmembrane domain, but no cytoplasmic death domain. Thi s receptor is not capable of inducing apoptosis, and is thought to function as an antagonistic receptor tha t protects cells from TRAIL-induced apoptosis. This gene was found to be a p53-regulated DNA damage-in ducible gene. The expression of this gene was detected in many normal tissues but not in most cancer cell lines, which may explain the specific sensitivity of cancer cells to the apoptosis-inducing activity of TRAIL .
------	--

基因ID	8794
基因名	TNFRSF10C
Swiss	O14798
别名	TNFRSF10C;CD263;DCR1;DCR1-TNFR;LIT;TRAIL-R3;TRAILR3;TRID

产品验证



Western blot analysis of TNFRSF10C expressed in Mouse liver,Rat liver using TNFRSF10C Rabbit pAb at 1 :1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn