

VHL Rabbit pAb

货号: B19987

产品信息

反应	Human, Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	18kDa/19kDa/24kDa
实测分子量	18kDa, 24kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse brain,Mouse kidney,MCF-7,Raji
细胞定位	Cytoplasm,Membrane,Nucleus,Nucleus,Peripheral membrane protein
纯化	Affinity purification

抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human VHL (NP_937799.1).
序列	MPRRAENWDEAEVGAEEAGVEEYGPEEDGGEESGAEESGPEESGPEELGAEEEMEAGRPRPVLRSVNSREPSQVIFCNR SPRVVLPVWLNFDGEPQPYPT


靶点信息

研究背景	Von Hippel-Lindau syndrome (VHL) is a dominantly inherited familial cancer syndrome predisposing to a
	variety of malignant and benign tumors. A germline mutation of this gene is the basis of familial inheritan
	ce of VHL syndrome. The protein encoded by this gene is a component of the protein complex that includ
	es elongin B, elongin C, and cullin-2, and possesses ubiquitin ligase E3 activity. This protein is involved in
	the ubiquitination and degradation of hypoxia-inducible-factor (HIF), which is a transcription factor that pl

ays a central role in the regulation of gene expression by oxygen. RNA polymerase II subunit POLR2G/RPB 7 is also reported to be a target of this protein. Alternatively spliced transcript variants encoding distinct i soforms have been observed.

基因ID	7428
基因名	VHL
Swiss	P40337
别名	VHL;HRCA1;RCA1;VHL1;pVHL;PVHL

产品验证



Western blot analysis of VHL expressed in Mouse brain, Mouse kidney, MCF-7 using VHL Rabbit pAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Bl ocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn