

STK36 Rabbit pAb

货号: B20003

产品信息

反应	Human, Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	141kDa/143kDa
实测分子量	144kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse brain, Mouse testis
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 250-450 of human STK3 6 (NP_056505.2).
序列	YHPFIAGHVTIITEPAGPDLGTPFTSRLPPELQVLKDEQAHRLAPKGNQSRILTQAYKRMAEEAMQKKHQNTGPALEQEDKT SKVAPGTAPLPRLGATPQESSLLAGILASELKSSWAKSGTGEVPSAPRENRTTPDCERAFPEERPEVLGQRSTDVVDLENE EPDSDNEWQHLLETTEPVPIQLKAPLTLLCNPDFCQRI

靶点信息

研究背景	This gene encodes a member of the serine/threonine kinase family of enzymes. This family member is si milar to a Drosophila protein that plays a key role in the Hedgehog signaling pathway. This human protein is a positive regulator of the GLI zinc-finger transcription factors. Knockout studies of the homologous mo use gene suggest that defects in this human gene may lead to congenital hydrocephalus, possibly due to a functional defect in motile cilia. Because Hedgehog signaling is frequently activated in certain kinds of g astrointestinal cancers, it has been suggested that this gene is a target for the treatment of these cancer s. Alternative splicing of this gene results in multiple transcript variants.
基因ID	27148
基因名	STK36
Swiss	Q9NRP7
别名	STK36;FU

产品验证



Western blot analysis of STK36 expressed in Mouse brain, Mouse testis using STK36 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn