

PRKCI Rabbit pAb

货号: B20010

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	68kDa
实测分子量	68kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-251MG,Mouse brain,Rat brain
细胞定位	
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-596 of human PRKCI (P41743).
序列	MPTQRDSSTMSHTVAGGGSGDHSHQVRVKAYRGDIMITHFEPSISFEGLCNEVRDMCSFDNEQLFTMKWIDEEGDPC TVSSQLELEAFRLYELNKDSELLIHVFPCVPERPGMPCPGEDKSIYRRGARRWRKLYCANGHTFQAKRFNRRAHCAICTDR IWGLGRQGYKCINCKLLVHKKCHKLVTIECGRHSPLQEPVMPMDQSSMHSDHAQTVIPYNPSSHESLDQVGEEKEAMNT RESGKASSSLGLQDFDLLRVIGRGSYAKVLLVRLKKTDRIYAMKVVKKELVNDDDEDIDWVQTEKHVFHQASNHPFVLGLH SCFQTESRLFFVIEVNNGGDLMFHMQRQRKLPEEHARFYSAEISLALNYLHERGIYRDLKLDNVLLDSEGHIKLTDYGMCK EGLRPGDTTSTFCGTPNYIAPEILRGEDYGFSDWWALGVLMFEMMAGRSPFDIVGSSDNPDQNTEDYLFQVILEKQIRIP RSLSVKAASVLKSFLNKDPKERLGCHPQTGFADIQGHFPRNVDWDMMEQKQVVPFKPNISGEFGLDNFDSQFTNEPV QLTPDDDDIVRKIDQSEFEGFEYINPLLMSAECV

靶点信息

研究背景	This gene encodes a member of the protein kinase C (PKC) family of serine/threonine protein kinases. The PKC family comprises at least eight members, which are differentially expressed and are involved in a wide variety of cellular processes. This protein kinase is calcium-independent and phospholipid-dependent. It is not activated by phorbol esters or diacylglycerol. This kinase can be recruited to vesicle tubular clusters (VTCs) by direct interaction with the small GTPase RAB2, where this kinase phosphorylates glyceraldehyde-3-phosphate dehydrogenase (GAPD/GAPDH) and plays a role in microtubule dynamics in the early secretory pathway. This kinase is found to be necessary for BCL-ABL-mediated resistance to drug-induced apoptosis and therefore protects leukemia cells against drug-induced apoptosis. There is a single exon pseudogene mapped on chromosome X.
基因ID	5584
基因名	PRKCI
Swiss	P41743
别名	PKCI;DXS1179E;nPKC-iota;PRKCI

产品验证

Western blot analysis of PRKCI expressed in U-251MG,Mouse brain,Rat brain using PRKCI Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



实验步骤

访问官网浏览详情: www.ablybio.cn