

GORASP2 Rabbit pAb

货号: B20104

产品信息

Human,Mouse,Rat
Rabbit
Polyclonal
WB IF/ICC
WB: 1:500 - 1:2000 F/ICC: 1:50 - 1:200
39kDa/47kDa/48kDa
55kDa
Liquid
Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Jnconjugated
.02,HeLa,Jurkat,MCF7,K-562,SW480,Mouse liver
Golgi apparatus membrane,Lipid-anchor,Membrane,Peripheral membrane protein
Affinity purification
γ · · · · · · · · · · · · · · · · · · ·

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 353-452 of human GOR ASP2 (NP_056345.3).
序列	PGIAPLPLPSEFLPSFPLVPESSSAASSGELLSSLPPTSNAPSDPATTTAKADAASSLTVDVTPPTAKAPTTVEDRVGDSTPV SEKPVSAAVDANASESP

靶点信息

研究背景	This gene encodes a member of the Golgi reassembly stacking protein family. These proteins may play a role in the stacking of Golgi cisternae and Golgi ribbon formation, as well as Golgi fragmentation during a poptosis or mitosis. The encoded protein also plays a role in the intracellular transport of transforming gro wth factor alpha and may function as a molecular chaperone. A pseudogene of this gene is located on the short arm of chromosome 2. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.
------	---

基因ID	26003
基因名	GORASP2
Swiss	Q9H8Y8
别名	GORASP2;GOLPH6;GRASP55;GRS2;p59

产品验证



Western blot analysis of GORASP2 expressed in Mouse liver,LO2,HeLa using GORASP2 Rabbit pAb at 1: 1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane . Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn