

# IGFALS Rabbit pAb

货号: **B20161**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	66kDa/70kDa
实测分子量	66kDa/72kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse lung,Rat liver
细胞定位	Secreted,extracellular space
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 396-605 of human IGFA LS (NP_004961.1).
序列	SCLGRIRPHTFTGLSGLRRLFLKDNGLVGIEEQLWGLAELLELDLTSNQLTHLPHRLFQGLGKLEYLLLSRNRLAELPADAL GPLQRAFWDVSHNRLEALPNSLLAPLGRRLRYLSLRNNSLRFTFPQPPGLERLWLEGNPWDCGCPKALRDFALQNPSAV PRFVQAICEGDDCQPPAYTYNNITCASPPPEVVGDLDRDLSEAHFAPC

## 靶点信息

研究背景	The protein encoded by this gene is a serum protein that binds insulin-like growth factors, increasing their half-life and their vascular localization. Production of the encoded protein, which contains twenty leucine-rich repeats, is stimulated by growth hormone. Defects in this gene are a cause of acid-labile subunit deficiency, which manifests itself in a delayed and slow puberty. Three transcript variants encoding two different isoforms have been found for this gene.
------	--

基因ID	3483
基因名	IGFALS
Swiss	P35858
别名	IGFALS;ACLSL;ALS

## 产品验证



Western blot analysis of IGFALS expressed in Mouse lung,Rat liver using IGFALS Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)