

LILRB2/CD85d/ILT4 Rabbit pAb

货号: B20194

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:1000 - 1:4000
理论分子量	51kDa/55kDa/64kDa/65kDa
实测分子量	65kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse liver,Mouse kidney,Mouse blood cell,Rat liver,Rat kidney,Rat blood cell
细胞定位	Membrane,Single-pass type I membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 22-240 of human LILRB2/CD85d/ILT4 (Q8N423).
序列	QTGTIPKPTLWAEPSVITQGSPVTLSCQGSLEAQEYRLYREKKSASWITRIRPELVKNGQFHIPSITWEHTGRYGCQYYSR ARWSELSDDLVLVMTGAYPKPTLSAQSPVVTSGGRVTLQCESQVAFGGFILCKEGEEEHPQCLNSQPHARGSSRAIFSV GPVSPNRRRWSHRCYGYDLNSPYVWSSPSDLLELLVPGVSKKPSLSVQPGPVVAPGES

靶点信息

研究背景	This gene is a member of the leukocyte immunoglobulin-like receptor (LIR) family, which is found in a gene cluster at chromosomal region 19q13.4. The encoded protein belongs to the subfamily B class of LIR receptors which contain two or four extracellular immunoglobulin domains, a transmembrane domain, and two to four cytoplasmic immunoreceptor tyrosine-based inhibitory motifs (ITIMs). The receptor is expressed on immune cells where it binds to MHC class I molecules on antigen-presenting cells and transduces a negative signal that inhibits stimulation of an immune response. It is thought to control inflammatory responses and cytotoxicity to help focus the immune response and limit autoreactivity. Multiple transcript variants encoding different isoforms have been found for this gene.
基因ID	10288
基因名	
Swiss	Q8N423
别名	LILRB2;CD85D;ILT-4;ILT4;LIR-2;LIR2;MIR-10;MIR10

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn