

LAIR1 Rabbit pAb

货号: B20202

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	23kDa/29kDa/31kDa
实测分子量	40-60KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	THP-1,HL-60,HeLa(Negative control),U-937
细胞定位	Cell membrane,Single-pass type I membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 22-165 of human LAIR1 (Q6GTX8).
序列	QEEDLPRPSISAEPGTVIPLGSHVTFVCRGPVGVQTFRLERESRSTYNDTEDVSQASPSESEARFRIDSVSEGNAGPYRCIY YKPPKWSEQSDYLELLVKETSGGPDSPDTEPGSSAGPTQRPSDNSHNEHAPASQGLKAEHLY

靶点信息

研究背景	The protein encoded by this gene is an inhibitory receptor found on peripheral mononuclear cells, includin g natural killer cells, T cells, and B cells. Inhibitory receptors regulate the immune response to prevent lys is of cells recognized as self. The gene is a member of both the immunoglobulin superfamily and the leuk ocyte-associated inhibitory receptor family. The gene maps to a region of 19q13.4 called the leukocyte re ceptor cluster, which contains at least 29 genes encoding leukocyte-expressed receptors of the immunogl obulin superfamily. The encoded protein has been identified as an anchor for tyrosine phosphatase SHP-1, and may induce cell death in myeloid leukemias. Alternative splicing results in multiple transcript varian ts.
基因ID	3903
基因名	LAIR1
Swiss	Q6GTX8
别名	LAIR1;CD305;LAIR-1

产品验证



Western blot analysis of LAIR1 expressed in THP-1,HL-60,HeLa(Negative control) using LAIR1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per la ne. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn