

CDCA8 Rabbit pAb

货号: B20285

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:100 - 1:200
理论分子量	31kDa
实测分子量	36-38kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HL-60,HeLa,MCF7,Mouse spleen
细胞定位	Chromosome,Cytoplasm,Nucleus,centromere,cytoskeleton,nucleolus,spindle
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-280 of human CDCA8 (Q53HL2).
序列	MAPRKGSSRVAKTNSLRRRKLASFLKDFDREVEIRIKQIESDRQNLLKEVDNLYNIEILRLPKALREMNWLDYFALGGNKQA LEEAAATADLDITEINKLTAEAIQTPLKSAKTRKVIQVDEMIVEEEEEENERKNLQTARVKRCPPSKKRTQSIQKGKGKRSS RANTVTPAVGRLEVSMVKPTPGLTPRFDSDRVFKTPGLRTPAAGERIYNISGNGSPLADSKEIFLTPVVGGSLSRLLASDLQ RHSIAQLDPEALGNIKKLSNRLAQICSSIRTHK

靶点信息

研究背景	This gene encodes a component of the chromosomal passenger complex. This complex is an essential re- gulator of mitosis and cell division. This protein is cell-cycle regulated and is required for chromatin-induc- ed microtubule stabilization and spindle formation. Alternate splicing results in multiple transcript variant s. Pseudogenes of this gene are found on chromosomes 7, 8 and 16.
------	---

基因ID	55143
基因名	CDCA8
Swiss	Q53HL2
别名	CDCA8;BOR;BOREALIN;DasraB;MESRGP;borealin

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn