

NR4A2 Rabbit pAb

货号**: B20472**

产品信息

| 反应 | Human, Mouse |
|-------|---|
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:1000 |
| 理论分子量 | 59kDa/66kDa |
| 实测分子量 | 67KDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HeLa,Jurkat,Raji,SH-SY5Y,Mouse brain |
| 细胞定位 | Cytoplasm,Nucleus |
| 纯化 | Affinity purification |

抗原信息


| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 90-170 of human NR4A 2 (NP_006177.1). | |
|------|---|--|
| 序列 | IKVEDIQMHNYQQHSHLPPQSEEMMPHSGSVYYKPSSPPTPTTPGFQVQHSPMWDDPGSLHNFHQNYVATTHMIEQRK TPV | |

靶点信息

| 研究背景 | This gene encodes a member of the steroid-thyroid hormone-retinoid receptor superfamily. The encoded protein may act as a transcription factor. Mutations in this gene have been associated with disorders relat ed to dopaminergic dysfunction, including Parkinson disease, schizophernia, and manic depression. Misre gulation of this gene may be associated with rheumatoid arthritis. Alternatively spliced transcript variants have been described, but their biological validity has not been determined. |
|--------------|---|
| 基因 ID | 4929 |

| 基因名 | NR4A2 |
|-------|----------------------------------|
| Swiss | P43354 |
| 别名 | NR4A2;HZF-3;NOT;NURR1;RNR1;TINUR |

产品验证



Western blot analysis of NR4A2 expressed in Mouse brain,HeLa,Jurkat using NR4A2 Rabbit pAb at 1:100 0. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blo cking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn