

Phospho-MAP3K5-S967 Rabbit pAb

货号: B20509

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	
实测分子量	155KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat
细胞定位	Cytoplasm,Endoplasmic reticulum
纯化	Affinity purification

抗原信息

抗原信息	A phospho specific peptide corresponding to residues surrounding S967 of human MAP3K5
序列	


靶点信息

研究背景

Mitogen-activated protein kinase (MAPK) signaling cascades include MAPK or extracellular signal-regulate d kinase (ERK), MAPK kinase (MKK or MEK), and MAPK kinase kinase (MAPKKK or MEKK). MAPKK kinase/ME KK phosphorylates and activates its downstream protein kinase, MAPK kinase/MEK, which in turn activate s MAPK. The kinases of these signaling cascades are highly conserved, and homologs exist in yeast, Dros ophila, and mammalian cells. MAPKKK5 contains 1,374 amino acids with all 11 kinase subdomains. North ern blot analysis shows that MAPKKK5 transcript is abundantly expressed in human heart and pancreas. T he MAPKKK5 protein phosphorylates and activates MKK4 (aliases SERK1, MAPKK4) in vitro, and activates c-Jun N-terminal kinase (JNK)/stress-activated protein kinase (SAPK) during transient expression in COS and 293 cells; MAPKKK5 does not activate MAPK/ERK.

基因ID	4217
基因名	MAP3K5
Swiss	Q99683
别名	MAP3K5;ASK1;MAPKKK5;MEKK5

产品验证



Western blot analysis of Phospho-MAP3K5-S967 expressed in Jurkat using Phospho-MAP3K5-S967 Rab bit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 3 0ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure tim e: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn