

BMPR2 Rabbit pAb

货号**: B20514**

产品信息

| 反应 | Human,Mouse,Rat |
|-------|---|
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB IHC |
| 推荐浓度 | WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 |
| 理论分子量 | 59kDa/115kDa |
| 实测分子量 | 100kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | mouse lung,rat kidney |
| 细胞定位 | Cell membrane,Single-pass type I membrane protein |
| 纯化 | Affinity purification |

抗原信息

| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 27-150 of human BMPR 2 (NP_001195.2). |
|------|---|
| 序列 | SQNQERLCAFKDPYQQDLGIGESRISHENGTILCSKGSTCYGLWEKSKGDINLVKQGCWSHIGDPQECHYEECVVTTTPP SIQNGTYRFCCCSTDLCNVNFTENFPPPDTTPLSPPHSFNRDET |

靶点信息

| 研究背景 | This gene encodes a member of the bone morphogenetic protein (BMP) receptor family of transmembran e serine/threonine kinases. The ligands of this receptor are BMPs, which are members of the TGF-beta su perfamily. BMPs are involved in endochondral bone formation and embryogenesis. These proteins transd uce their signals through the formation of heteromeric complexes of two different types of serine (threoni ne) kinase receptors: type I receptors of about 50-55 kD and type II receptors of about 70-80 kD. Type II r eceptors bind ligands in the absence of type I receptors, but they require their respective type I receptors for signaling, whereas type I receptors require their respective type II receptors for ligand binding. Mutati ons in this gene have been associated with primary pulmonary hypertension, both familial and fenflurami ne-associated, and with pulmonary venoocclusive disease. |
|-------|--|
| 基因ID | 659 |
| 基因名 | BMPR2 |
| Swiss | Q13873 |
| 别名 | BMPR2;BMPR-II;BMPR3;BMR2;BRK-3;POVD1;PPH1;T-ALK |

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn