

HOXD8 Rabbit pAb

货号**: B20556**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	
实测分子量	32kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SKOV3,Mouse ovary,Rat ovary
细胞定位	nucleus
纯化	Affinity purification

抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 200 to the C-terminus of human HOX D8 (NP_001186675.1).	
序列	RQTYSRFQTLELEKEFLFNPYLTRKRRIEVSHALALTERQVKIWFQNRRMKWKKENNKDKFPVSRQEVKDGETKKEAQELE EDRAEGLTN	

靶点信息

研究背景	This gene belongs to the homeobox family of genes. The homeobox genes encode a highly conserved fa mily of transcription factors that play an important role in morphogenesis in all multicellular organisms. M ammals possess four similar homeobox gene clusters, HOXA, HOXB, HOXC and HOXD, located on differen t chromosomes, consisting of 9 to 11 genes arranged in tandem. This gene is one of several homeobox H OXD genes located in a cluster on chromosome 2. Deletions that remove the entire HOXD gene cluster or the 5' end of this cluster have been associated with severe limb and genital abnormalities. In addition to effects during embryogenesis, this particular gene may also play a role in adult urogenital tract function. Alternate splicing results in multiple transcript variants.
基因ID	3234
基因名	HOXD8
Swiss	P13378
别名	HOXD8;HOX4;HOX4E;HOX5.4

产品验证



Western blot analysis of HOXD8 expressed in SKOV3,Mouse ovary,Rat ovary using HOXD8 Rabbit pAb a t 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per la ne. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn