

Histone H4 Rabbit pAb

货号**: B20626**

产品信息

反应	Human,Mouse,Rat,Other (Wide Range)
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	11kDa
实测分子量	11kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,NIH/3T3,C6
细胞定位	Chromosome,Nucleus
纯化	Affinity purification

抗原信息

抗原信息	A synthetic monomethylated peptide around R3 of human Histone H4检索 (NP_003529.1).
序列	SGRGK

靶点信息

研究背景	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octa mer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The I inker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chr omatin into higher order structures. This gene is intronless and encodes a replication-dependent histone t hat is a member of the histone H4 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6.

基因ID	8359
基因名	
Swiss	P62805
别名	H4C2; H4C3; H4C4; H4C5; H4C6; H4C8; H4C9; H4FA; H4-16; H4C11; H4C12; H4C13; H4C14; H4C15; HIST 1H4A

产品验证



Western blot analysis of Histone H4 expressed in HeLa,NIH/3T3,C6 using Histone H4 Rabbit pAb at 1: 1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per Ian e. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn