

MRPS7 Rabbit pAb

货号: B20693

产品信息

| 反应 | Human, Mouse, Rat |
|-------|---|
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB IF/ICC |
| 推荐浓度 | WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 |
| 理论分子量 | 28kDa |
| 实测分子量 | 28kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HepG2,A-549,HT-29,Mouse liver,Mouse kidney,Rat liver,Rat kidney |
| 细胞定位 | mitochondrial inner membrane |
| 纯化 | Affinity purification |


抗原信息

| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 110-190 of human MRP S7 (NP_057055.2). |
|------|--|
| 序列 | LMIQTLEAVKRKQFEKYHAASAEEQATIERNPYTIFHQALKNCEPMIGLVPILKGGRFYQVPVPLPDRRRRFLAMKWMITE |

靶点信息

| 研究背景 | Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a l arge 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribo somes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryo tic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the m itoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy rec ognition by sequence homology. This gene encodes a 28S subunit protein. In the prokaryotic ribosome, the comparable protein is thought to play an essential role in organizing the 3' domain of the 16 S rRNA in the vicinity of the P- and A-sites. Pseudogenes corresponding to this gene are found on chromosomes 8p and 12p. |
|--------------|--|
| 基因 ID | 51081 |
| 基因名 | MRPS7 |
| Swiss | Q9Y2R9 |
| 别名 | MRPS7;MRP-S;RP-S7;RPMS7;S7mt;bMRP27a |

产品验证



Western blot analysis of MRPS7 expressed in Mouse liver,Rat liver,HepG2 using MRPS7 Rabbit pAb at 1: 1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn