

ZCCHC3 Rabbit pAb

货号**: B20738**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	43kDa
实测分子量	43kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	LO2,mouse liver
细胞定位	cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 215-335 of human ZCC HC3 (NP_149080.2).
序列	SFRSAEKLALFLRVYEEKREQEDCWENFVVLGRSKSSLKTLFILFRNETVDVEDIVTWLKRHCDVLAVPVKVTDRFGIWTGE YKCEIELRQGEGGVRHLPGAFFLGAERGYSWYKGQPKTC

靶点信息

研究背景	Nucleic acid-binding protein involved in innate immune response to DNA and RNA viruses. Binds DNA and RNA in the cytoplasm and acts by promoting recognition of viral nucleic acids by virus sensors, such as D DX58/RIG-I, IFIH1/MDA5 and CGAS. Acts as a co-sensor for recognition of double-stranded DNA (dsDNA by cGAS in the cytoplasm, thereby playing a role in innate immune response to cytosolic dsDNA and DNA vir us. Binds dsDNA and probably acts by promoting sensing of dsDNA by CGAS, leading to enhance CGAS oli gomerization and activation. Promotes sensing of viral RNA by RIG-I-like receptors proteins DDX58/RIG-I a nd IFIH1/MDA5 via two mechanisms: binds double-stranded RNA (dsRNA, enhancing the binding of DDX58 /RIG-I and IFIH1/MDA5 to dsRNA and promotes 'Lys-63'-linked ubiquitination and subsequent activation of DDX58/RIG-I and IFIH1/MDA5.
基因 ID	85364
基因名	ZCCHC3
Swiss	Q9NUD5
别名	ZCCHC3;C20orf99

产品验证



Western blot analysis of ZCCHC3 expressed in LO2,mouse liver using ZCCHC3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buff er: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn