

RASD2 Rabbit pAb

货号: B20790

产品信息

反应	Human, Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	30kDa
实测分子量	30kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse lung
细胞定位	plasma membrane,synapse
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 135-263 of human RAS D2 (NP_055125.2).
序列	MVICGNKNDHGELCRQVPTTEAELLVSGDENCAYFEVSAKKNTNVDEMFYVLFSMAKLPHEMSPALHRKISVQYGDAFHP RPFCMRRVKEMDAYGMVSPFARRPSVNSDLKYIKAKVLREGQARERDKC

靶点信息

研究背景	This gene belongs to the Ras superfamily of small GTPases and is enriched in the striatum. The encoded protein functions as an E3 ligase for attachment of small ubiquitin-like modifier (SUMO). This protein also binds to mutant huntingtin (mHtt), the protein mutated in Huntington disease (HD). Sumoylation of mHTT by this protein may cause degeneration of the striatum. The protein functions as an activator of mechanis tic target of rapamycin 1 (mTOR1), which in turn plays a role in myelination, axon growth and regeneratio n. Reduced levels of mRNA expressed by this gene were found in HD patients.
------	---

基因ID	23551
基因名	RASD2
Swiss	Q96D21
别名	Rhes;TEM2;MGC:4834;RASD2

产品验证



Western blot analysis of RASD2 expressed in Mouse lung using RASD2 Rabbit pAb at 1:1000. Secondary a ntibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn